

On the current status of open-cluster parameters

E. Paunzen, M. Netopil

Institut für Astronomie der Universität Wien, Türkenschanzstr. 17, 1180 Wien, Austria

4 February 2008

ABSTRACT

We aim to characterize the current status of knowledge on the accuracy of open-cluster parameters such as the age, reddening and distance. These astrophysical quantities are often used to study the global characteristics of the Milky Way down to very local stellar phenomena. In general, the errors of these quantities are neglected or set to some kind of heuristic standard value. We attempt to give some realistic estimates for the accuracy of available cluster parameters by using the independent derived values published in the literature. In total, 6437 individual estimates for 395 open clusters were used in our statistical analysis. We discuss the error sources depending on theoretical as well as observational methods and compare our results with those parameters listed in the widely used catalogue by Dias et al. (2002). In addition, we establish a list of 72 open clusters with the most accurate known parameters which should serve as a standard table in the future for testing isochrones and stellar models.

Key words: Open clusters and associations: general

1 INTRODUCTION

The study of open clusters naturally introduces many advances, because they are physically related groups of stars held together by mutual gravitational attraction that were formed at roughly the same time from one large cosmic gas and dust cloud. Their evolutionary stages range from clouds where star formation still takes place at this moment to very old aggregates with turn-off points as late as solar type stars. Therefore, they represent samples of Population I stars of constant age and comparable intrinsic chemical composition, best suited to study processes related to stellar evolution and formation, and to fix lines or loci in several most important astrophysical diagrams. These procedures are statistical methods, for example fitting isochrones to determine the age, reddening, and distance of an open cluster, independent of individual peculiarities of members.

As the open clusters drift along their orbits, they are excellent tracers for the global kinematics and dynamical characteristics of the Milky Way itself. However, some of their members escape the cluster, due to velocity changes in mutual close encounters or tidal forces in the galactic gravitational field. The escaped stars continue to orbit on their own as field stars.

As a summary it can be concluded that open clusters allow the study of a significant number of astrophysical parameters and models from the global scale of the Milky Way down to processes in individual stars from the Pre-Main-Sequence to Post-AGB evolution.

But what about the accuracy of the presently known

open-cluster parameters? Surprisingly enough, there is, to our knowledge, no work within the last ten years which deals with this question in detail. Although there are several new catalogues (Dias et al. 2002 and Kharchenko et al. 2005) published and a flood of new parameter estimates, mainly because the common use of CCDs permits the observation of fainter and more distant clusters, there is no published statistical analysis describing if and how the accuracy has improved.

If we look at recent statistical papers dealing with open-cluster parameters (e.g. Chen, Hou & Wang 2003, Dias & Lépine 2005), none of them have any detailed error treatment included. Although apparent “mean values” for the age, distance and metallicity are used, the level of accuracy is completely ignored in the final conclusions.

In this work we try to fill this gap by presenting a detailed analysis of the current available open-cluster parameters and the corresponding errors. For this purpose we have searched the literature up to November 2005 for individual cluster parameter estimates. The primary goal is to establish a list of “standard” open clusters covering a wide range of ages, reddenings and distances from the Sun selected on the basis of the smallest errors from the available parameters in the literature. These clusters should serve as a standard set for future investigations. We also compare the derived mean values with those of the most recent catalogue by Dias et al. (2002) resulting in an overall satisfactory agreement.




Figure 1. The errors of the mean for the distance, reddening and age versus the number of available estimates in the literature for the complete sample of 395 open clusters. They are smaller for the distance and reddening if more observations are available, but the mean errors for the age estimate are more or less constant. There are only few open clusters with more than 10 measurements which are not included in these diagrams to avoid a bias due to poor number statistics.

2 DATA SELECTION AND PREPARATION

For the determination of mean values for the age, reddening and distance, we have used the following data sources: Janes, Tilley & Lyngå, (1988), Malysheva (1997), Dambis (1999) Dutra & Bica (2000), Loktin et al. (2001), Tadross (2001), Lata et al. (2002, and references therein), Kharchenko et al. (2005) and an updated list of individual publications as listed in WEBDA (http://www.univie.ac.at/webda/recent_data.html) which includes more than 300 different papers. The latter was mainly performed within the framework of ADS (<http://cdsads.u-strasbg.fr/>). The individual references were further checked for sources from the literature to guarantee a maximum of available estimates accessible to the community. All selected publications list $E(B - V)$, $\log t$ and the distance from the Sun. The only exception is Malysheva (1997) who lists $E(b - y)$ which was converted by a factor of 1.43 to $E(B - V)$. For the further analysis we have used $A_V = 3.1E(B - V)$ which is the best mean value for most regions in the Milky Way (Winkler 1997).

We have carefully checked whether the listed parameters for each open cluster in the individual references are in-

dependent of each other and not used by two authors twice. Kharchenko et al. (2005), for example, used data by Loktin et al. (2001). Such duplicated data were not used in the subsequent analysis.

Janes et al. (1988) studied all photometric measurements available at that time to them and introduced weights to derive the cluster parameters. They summarize the knowledge and results for the given time and supersede the catalogue by Lyngå (1987).

The catalogue of Dias et al. (2002) was not used at this stage because it already lists averaged values from the literature. Furthermore we made a comprehensive comparison with the results of this work (Sect. 4).

The metallicity is also a free parameter when fitting isochrones to photometric data. Chen et al. (2003) compiled a catalogue of metallicities and analysed this sample. They derived a radial iron gradient of $-0.063(8)$ dex kpc^{-1} for the Milky Way with an intrinsic spread of the overall open cluster metallicities of ± 0.2 dex. This is in excellent agreement with the scatter of the metallicity found for F and G type stars in the solar neighbourhood (Karatas, Bilir & Schuster 2005). The effect of different metallicities to the isochrone fitting procedure is extensively discussed by Pinsonneault et al. (2004). The ΔM_V values range from $+0.1$ to -0.3 mag for $[\text{Fe}/\text{H}] = \pm 0.2$ dex and constant effective temperature sensitive color indices. Taking a difference of 0.3 mag for the distance modulus introduced by an incorrect treatment of the metallicity, the error for 1 kpc is 150 pc or 15%. However, these discrepancies are only valid for effective temperatures cooler than 8000 K. For hotter temperatures, the difference in the metallicities result mainly in a shift of the temperature sensitive color indices (Schaerer et al. 1993). Lebreton (2000) presented a comparison of galactic field stars with known metallicities as well as accurate parallax measurements and appropriate isochrones (see Figure 6 therein). There is an apparent mismatch at any level of metallicity. However, the most significant differences are at large underabundances (> -2.5 dex) which are normally not found in open clusters of the Milky Way. There are only a few independent estimates of the metallicity in the literature. Twarog, Ashman & Anthony-Twarog (1997), Gratton (2000) and Chen et al. (2003) give a summary of the current knowledge with an error estimate of the individual values. Since then very few new estimates have been published. We have therefore not made a new and redundant error analysis of the metallicities for open clusters.

All individual data were considered as completely equal and no weights were attributed. No assessment of the corresponding reference for the used methods, isochrones and so on was done. This should guarantee that no unknown bias is introduced in the data set. We have rejected NGC 2451, which recent studies have shown that is formed by the superposition of two clusters at different distances, and Basel 11, which was wrongly assigned in the original survey. Therefore the former parameters determined for the joined data are no longer significant. Furthermore, Pena & Peniche (1994) reported significant inconsistencies for the analysis of NGC 7209 which we also neglected.

We have also not included the given “intrinsic” errors of individual references. The simple reason is that most authors list only an overall error (if any) for all measurements which is highly unrealistic and would introduce an unknown bias.




Figure 2. The distribution of the mean absolute errors of the distance and age as well as the standard deviation of the reddening for the complete sample of 395 open clusters with more than three independent measurements from the literature (Section 3). This figure summarizes the current accuracy of open-cluster parameters.

For the final list, we have averaged the data of open clusters for which at least three independent estimates of the age, reddening and distance are available. It contains 6437 individual estimates for 395 open clusters. As the last step, the standard deviations were calculated.

3 STATISTICAL ANALYSIS OF THE SAMPLE

Fitting isochrones to photometric data is not straightforward. The goodness of the fit depends on the following characteristics of the colour-magnitude diagram of an open cluster:

- Number of members
- Percentage of contamination by non-members
- Age, i.e. visibility of the turn-off point, an important source of errors for young open clusters
- Presence of spectroscopic binaries which shifts the main sequence by a maximum of 0.754 mag
- Presence of a red giant clump which makes an isochrone fitting much more significant
- Differential reddening




Figure 3. Distribution of the differences of the distance, age (both in units of the standard deviation of the mean) and reddening from the most recent catalogue by Dias et al. (2002) and the estimate from this work. The values of the Gaussian fits are shown in the panels.

- Available photometric indices, $(U - B)$ is, for example, an excellent indicator for the reddening and membership
- Quality of the data
- Metallicity

In addition, there is also a “human factor” which enters the estimate of the best fit to observational data. Andersen & Nordström (1999) show how different research groups interpret the same data set quite differently, using the excellent example of NGC 3680 (Figure 3 therein). This is not because one set of parameters is better than the other but the technique of isochrone fitting itself is not unambiguous. For this reason, very sophisticated computer algorithms were developed to overcome this problem (Jørgensen & Lindegren 2005). The method is rather simple: the locations of the color-magnitude diagrams are divided into boxes and the incidence of members within each box is counted. Those numbers are compared with the same parameterization of standard isochrones. Loktin et al. (2001) and Kharchenko et al. (2005), for example, used such a black box to derive cluster parameters. However, Jørgensen & Lindegren (2005) concluded that the systematic errors of computer algorithms are at least on the same level as those introduced by the human factor.

Another error source is the choice of appropriate isochrones. Grocholski & Sarajedini (2003) discussed the apparent differences of the five most commonly used sets of isochrones. They concluded that none of the theoretical isochrones reproduce the observational data in a consistent manner over the magnitude and colour range of the zero age main sequence. In particular, there are significant zero point and shape differences between the models and the observations. There are three reasons for these discrepancies: 1) the input physics has to be improved; 2) The transformation between theoretical and observational parameters; and 3) more standard open clusters have to be established for testing new models.

For our statistical analysis we have chosen the following parameters:

- $\sigma(D)/D$: the absolute error of the mean distances
- $\sigma E(B - V)$: the standard deviation of the mean reddening values
- $\sigma(\text{age})/\text{age}$: the absolute error of the mean ages (in units of Myrs and not in the logarithmic scale)

The parameters themselves are, in general, *not* independent from each other. For example, the distance depends on the total absorption.

Figure 1 shows the errors of the mean for the distance, reddening and age versus the number of available estimates in the literature for the complete sample of 395 open clusters. The errors of the means for the distance and reddening decrease with more available observations while the mean errors for the age estimate are more or less constant over the whole range. Note that there are also a few parameters which are estimated more than ten times, but these few data are not shown to avoid any poor number statistics.

The distribution of the different error estimates (Fig. 2) gives an idea of the current accuracy for the open-cluster parameters. The reddening is best known mainly because the main sequence can be fitted more easily for almost all open clusters. For about 90% of all clusters, the error is below 0.1 mag. The distances are also rather well determined: for about 80% of all aggregates, the absolute error is less than 20%. As expected, the estimate of the age of open clusters is the most difficult and introduces large uncertainties. Only 11% of the investigated open clusters have errors less than 20%. Even more than 30% exhibit an absolute error larger than 50% with extreme values of more than 200%. These characteristics are most certainly not due to wrong estimates or typographical errors in the reference but an intrinsic problem of the method itself.

How should one overcome the problem of a proper age estimate for open clusters? We have shown that the knowledge of the reddening and determining distances is satisfactory. If we look at the metallicity, the situation is rather poor. Most authors use solar abundances (e.g. Dutra & Bica 2000, Loktin et al. 2001 and Kharchenko et al. 2005) throughout the isochrone fitting process, this procedure is in most cases justified, but normally photometric data will not allow us do it any better. We believe that the only way out of the dilemma is to establish “standard” open clusters covering a wide range of distances, ages and reddening values which should allow, for example the testing of isochrones.

4 COMPARISON WITH THE CATALOGUE BY DIAS AND COWORKERS

The current best approximation, many authors use is the list of parameters published by Dias et al. (2002). We have used the open-cluster parameters of the October 2005 version of the catalogue available at <http://www.astro.iag.usp.br/~wilton/> for comparison with our results. The motivation for this comparison is to have an estimate of the statistical error introduced when using “mean” catalogue parameters instead of performing an independent analysis of its own.

We are aware that the catalogue of Dias et al. is a kind of compendium of different sources from the literature. They do not perform any weighting or averaging in the tradition of Lyngå. Our analysis, on the other hand, is based on averaged values from widely different methods and authors. It is therefore very interesting to examine if an independent check of this catalogue can confirm its validity.

For the analysis we have calculated the differences of the open-cluster parameters of the catalogue by Dias et al. (2002) and our mean estimate for the sample of 395 open clusters. For the reddening we have just used the differences as they were calculated. Otherwise, those difference were normalized in units of the standard deviation of the mean for the distance and age. The interpretation of those parameters is the following: for values between -1 and $+1$, the error of averaging the literature parameter is statistically at the same level as the uncertainty introduced by using those catalogued ones. Fitting a Gaussian distribution to the complete sample of values should result in a width of less than $2\sqrt{2\ln 2} \approx 2.355$ if both “error sources” have the same statistical significance and can therefore not be disentangled (Rees 1987).

Figure 3 shows the distribution of the differences of the distance, reddening and age as well as the basic parameters of a Gaussian fit to these distributions. The widths of all three distributions are in the expected range taking into account the overall statistical errors on the cluster parameters (Fig. 2). We therefore conclude that if one uses the parameters of the catalogue by Dias et al. (2002) the expected errors are comparable to those which we derived for averaging the independent values from the literature, a result which is in favor of the parameters from the mentioned catalogue from a statistical point of view.

5 A SET OF STANDARD CLUSTERS


If one aims at a statistical study of a group of clusters, a standard set for comparison is necessary. Mermilliod (1981), for example used the following clusters to establish empirical isochronous curves: Alpha Persei, Hyades, IC 4665, NGC 457, NGC 884, NGC 2281, NGC 2287, NGC 2362, NGC 2516, NGC 3532, NGC 3766, NGC 6231, NGC 6475 and the Pleiades. Those clusters still serve as references when studying red giants (Eigenbrod et al. 2004). Another interesting work in this respect is the one by Piatti, Claria & Bica (1998) who presented composite M_V versus $(V - I)_0$ diagrams for the following “standard” open clusters: IC 4651, NGC 3680, NGC 3766, NGC 6025, NGC 6067, NGC 6231, NGC 6242, NGC 6259, NGC 6451 and NGC 6633. They se-

Table 1. List of 72 suggested standard open clusters, sorted after the galactic longitude, from the literature. The listed parameters are calculated mean values with their standard deviations and should be treated in a statistical sense. The columns “*No*” denote the number of used independent estimates.

Cluster	Name	l	b	D [pc]	$\sigma(D)$ [pc]	No	$E(B - V)$ [mag]	$\sigma E(B - V)$ [mag]	No	age [Myr]	$\sigma(age)$ [Myr]	No
Collinder 394	C1850–204	14.88	−9.24	648	41	4	0.25	0.01	5	74	6	5
Berkeley 81	C1859–005	34.51	−2.07	2798	508	4	1.00	0.01	4	1000	1	4
IC 4756	C1836+054	36.38	+5.24	415	46	5	0.19	0.01	6	674	128	5
NGC 6834	C1950+292	65.70	+1.19	2147	59	4	0.73	0.03	4	65	18	4
NGC 6791	C1919+377	69.96	+10.90	4418	743	7	0.17	0.04	7	7850	2026	7
NGC 6871	C2004+356	72.64	+2.05	1675	132	7	0.46	0.03	8	9	2	8
NGC 6819	C1939+400	73.98	+8.48	2188	255	5	0.23	0.08	5	2172	442	5
NGC 6811	C1936+464	79.21	+12.02	1146	112	5	0.14	0.03	6	635	138	6
NGC 7044	C2111+422	85.89	−4.15	3097	145	6	0.63	0.06	6	1824	361	6
NGC 6939	C2030+604	95.90	+12.30	1321	274	6	0.45	0.08	6	1639	452	6
King 10	C2252+589	108.48	−0.40	3373	167	4	1.15	0.01	4	45	11	4
NGC 7789	C2354+564	115.53	−5.39	1924	182	8	0.25	0.02	9	1519	202	8
Berkeley 99	C2319+714	115.95	+10.11	4902	51	4	0.30	0.01	4	3162	1	4
King 12	C2350+616	116.12	−0.13	2453	124	5	0.58	0.03	6	11	1	5
Berkeley 2	C0022+601	119.71	−2.31	5129	383	4	0.80	0.01	4	794	1	4
NGC 129	C0027+599	120.27	−2.54	1632	56	8	0.56	0.03	9	62	15	9
Trumpler 1	C0132+610	128.22	−1.13	2356	511	9	0.57	0.04	9	30	6	9
Berkeley 64	C0217+656	131.92	+4.61	3860	229	5	1.05	0.01	5	1000	0.01	5
NGC 744	C0155+552	132.40	−6.16	1214	176	3	0.41	0.05	5	184	49	5
Melotte 20	Alpha Per	146.57	−5.86	173	12	8	0.09	0.03	9	43	18	9
NGC 1245	C0311+470	146.65	−8.93	2593	299	9	0.27	0.03	10	1065	276	10
King 7	C0355+516	149.77	−1.02	2211	200	5	1.27	0.05	5	603	121	5
NGC 1798	C0508+475	160.70	+4.85	4160	232	4	0.51	0.01	4	1421	16	4
Berkeley 18	C0518+453	163.63	+5.02	4772	723	4	0.47	0.01	4	4271	241	4
Melotte 22	Pleiades	166.57	−23.52	133	9	8	0.05	0.01	6	79	52	5
NGC 1778	C0504+369	168.90	−2.02	1560	105	5	0.32	0.04	7	129	29	7
NGC 2192	C0611+398	173.41	+10.65	3467	72	4	0.21	0.01	4	1072	48	4
Berkeley 69	C0521+326	174.43	−1.79	3121	311	4	0.67	0.05	4	867	48	4
NGC 2281	C0645+411	174.90	+16.88	488	48	4	0.08	0.02	6	431	105	6
Berkeley 17	C0517+305	175.65	−3.65	2575	88	4	0.63	0.05	4	9448	1776	4
Melotte 25	Hyades	180.06	−22.34	42	3	5	0.00	0.01	6	708	136	6
NGC 1647	C0443+189	180.34	−16.77	492	69	3	0.36	0.03	5	130	25	5
NGC 2158	C0604+241	186.63	+1.78	4403	589	5	0.44	0.06	5	1710	424	5
NGC 2266	C0640+270	187.79	+10.29	3490	180	4	0.10	0.01	4	736	77	4
NGC 2355	C0714+138	203.39	+11.80	2086	163	5	0.14	0.06	5	833	137	5
NGC 2632	Praesepe	205.92	+32.48	171	12	6	0.00	0.01	8	753	201	8
Berkeley 32	C0655+065	207.95	+4.40	3491	401	5	0.15	0.03	5	3477	698	5
NGC 2682	C0847+120	215.70	+31.90	820	47	12	0.05	0.02	13	4093	958	12
Melotte 111	Coma Ber	221.35	+84.03	86	7	5	0.00	0.01	6	522	82	6
Berkeley 39	C0744–044	223.46	+10.10	4283	425	4	0.12	0.01	4	7331	906	4
NGC 2548	C0811–056	227.87	+15.39	727	57	6	0.04	0.02	8	364	102	8
NGC 2506	C0757–106	230.56	+9.94	3315	219	8	0.06	0.04	9	1648	485	9
NGC 2539	C0808–126	233.71	+11.11	1270	162	5	0.08	0.02	7	490	129	6
NGC 2527	C0803–280	246.09	+1.86	581	29	4	0.07	0.03	5	619	163	5
NGC 2477	C0750–384	253.56	−5.84	1227	166	9	0.26	0.08	10	875	238	10
Melotte 66	C0724–476	259.56	−14.24	3784	828	5	0.16	0.01	5	4404	1145	5
Trumpler 10	C0846–423	262.79	+0.67	422	34	6	0.05	0.01	7	37	9	7
NGC 2660	C0840–469	265.93	−3.01	2859	71	5	0.37	0.03	5	1351	291	5
IC 2395	C0839–480	266.64	−3.59	792	131	4	0.09	0.02	5	15	4	5
NGC 3105	C0959–545	279.92	+0.26	7137	1710	6	1.04	0.06	6	21	3	5
IC 2581	C1025–573	284.59	+0.04	2215	317	5	0.42	0.02	6	13	3	6
Bochum 10	C1040–588	287.02	−0.31	2472	551	4	0.33	0.03	5	8	2	4
NGC 3532	C1104–584	289.57	+1.35	492	8	5	0.04	0.01	6	262	46	5
IC 2714	C1115–624	292.40	−1.80	1217	72	5	0.40	0.06	6	253	66	6
Melotte 105	C1117–632	292.90	−2.41	2094	159	7	0.48	0.05	7	224	53	7
Stock 14	C1141–622	295.22	−0.66	2439	326	4	0.24	0.02	5	10	2	5
Collinder 261	C1234–682	301.68	−5.53	2241	100	4	0.28	0.02	4	8812	201	4
ESO 096–SC04		305.35	−3.17	10402	1898	5	0.69	0.06	5	752	59	5

Table 1. continued.

Cluster	Name	l	b	D [pc]	$\sigma(D)$ [pc]	No	$E(B - V)$ [mag]	$\sigma E(B - V)$ [mag]	No	age [Myr]	$\sigma(age)$ [Myr]	No
NGC 5316	C1350–616	310.23	+0.12	1050	228	4	0.28	0.07	5	166	33	5
Pismis 19	C1426–607	314.71	−0.30	2149	435	4	1.46	0.01	4	973	127	4
NGC 5662	C1431–563	316.94	+3.39	684	60	7	0.32	0.01	8	77	20	8
NGC 5822	C1501–541	321.57	+3.59	787	75	6	0.14	0.06	7	913	246	7
NGC 6025	C1559–603	324.55	−5.88	725	94	5	0.16	0.01	6	74	22	6
NGC 6005	C1551–572	325.78	−2.99	2600	221	4	0.48	0.06	4	1100	204	4
NGC 6087	C1614–577	327.73	−5.43	893	57	7	0.20	0.03	9	78	19	9
NGC 6067	C1609–540	329.74	−2.20	1676	202	7	0.36	0.03	8	95	23	8
Lyngå 6	C1601–517	330.37	+0.32	1792	272	5	1.33	0.04	5	35	9	4
NGC 6208	C1645–537	333.76	−5.76	1096	170	5	0.19	0.02	6	1354	368	6
NGC 6253	C1655–526	335.46	−6.25	1567	108	7	0.25	0.08	7	3949	1086	7
NGC 6250	C1654–457	340.68	−1.92	936	56	5	0.37	0.02	6	22	5	5
NGC 6396	C1734–349	353.93	−1.77	1276	117	5	0.95	0.02	5	30	8	4
NGC 6405	C1736–321	356.58	−0.78	473	16	5	0.14	0.02	7	71	21	7

**Figure 4.** Distribution of the distance, reddening and age of the 72 suggested standard clusters as listed in Table 1.

lected these clusters because of accurately determined fundamental parameters. The final diagrams are used as a homogeneous set of empirical isochrones in the age range between 5 Myr and 4 Gyr for the study of faint reddened open clusters. The lists of Mermilliod (1981) and Piatti et al. (1998) only coincide in three cases: NGC 3766, NGC 6231

and NGC 6475. This already indicates that a common set of standard open clusters is very much needed.

The lists of standard clusters given in the above mentioned references have been chosen based on the available photometric, spectroscopic and kinematical data. However, as an example, Paunzen & Maitzen (2002) have shown that the photometric data of NGC 6451 are not intrinsically consistent and suffer from an offset of one magnitude in Johnson V . Our approach is different and not based on the current available data for individual open clusters. On the contrary, our list should trigger new observations to further improve the knowledge of the listed cluster parameters. We present a list of open clusters with a significant number of independent investigations and parameters with small errors.

We have chosen the rather strict criteria (especially for the age estimate) for the errors of the parameters:

- $\sigma(D)/D \leq 25\%$
- $\sigma E(B - V) \leq 0.1 \text{ mag}$
- $\sigma(age)/age \leq 30\%$

In total, 72 open clusters, listed in Table 1, fulfill these criteria. Figure 4 shows the distribution of the distances, reddening and ages for these proposed standard clusters. They cover the complete range of known cluster parameters within the Milky Way. These open clusters should serve as observational test cases for further theoretical developments. A closer inspection of these open clusters in WEBDA reveals that, for example, the data of NGC 6208 and NGC 6811 are still very unsatisfactory which calls for further high precision observational investigations.

6 CONCLUSIONS

Using open clusters for studying global and local phenomena is very much dependent on the accuracy of the used cluster parameters such as the age, reddening and distance. This is also vital for testing theoretical models. We have made a statistical study of the published cluster parameters summarizing the results for 395 open clusters. While the reddening seems to be quite accurate, the ages and distances suffer from severe uncertainties. For about 90% of all

studied clusters, the error of the reddening is below 0.1 mag. For the distance we find absolute errors of less than 20% for about 80% of the aggregates. However, only 11% of the investigated open clusters have errors of the ages which are less than 20%; there are extreme discrepancies of more than 200%. This calls for a homogeneous set of isochrones together with a solid fitting technique.

We have compared the errors of the mean cluster parameters with the differences of those listed in the open cluster catalogue by Dias et al. (2002) which is widely used. Both error distributions have the same statistical significance. Using the parameters from the mentioned catalogue, therefore, introduces the same error level as using averaged values from the literature, an important result when using this catalogue.

As last step, a set of 72 open clusters with the most accurate (in a statistical sense) parameters were established. Those clusters cover a wide range of reddenings, ages as well as distances and should serve in the future as a standard set for further observational efforts and for testing theoretical models.

Acknowledgements. We thank our referee, Jean-Claude Mermilliod, for very useful comments which helped to improve this manuscript significantly. This research was performed within the projects *P17580* and *P17920* of the Austrian Fonds zur Förderung der wissenschaftlichen Forschung (FwF). M. Netopil acknowledges the support by a “Forschungsstipendium” of the University of Vienna. Use was made of the SIMBAD database, operated at the CDS, Strasbourg, the WEBDA database, operated at the University of Vienna and the NASA Astrophysics Data System.

REFERENCES

- Andersen J., Nordström B., 1999, in Giménez A., Guinan E.F., Montesinos B., eds, ASP Conf. Ser. 173, Theory and Tests of Convection in Stellar Structure, San Francisco, p. 31
- Chen L., Hou J.L., Wang J.J., 2003, AJ, 125, 1397
- Dambis A.K., 1999, Astronomy Letters, 25, 7
- Dias W.S., Lépine J.R.D., 2005, ApJ, 629, 825
- Dias W.S., Alessi B.S., Moitinho A., Lépine J.R.D., 2002, A&A, 389, 871
- Dutra C.M., Bica E., 2000, A&A, 359, 347
- Eigenbrod A., Mermilliod J.-C., Clariá J.J., Andersen J., Mayor M., 2004, A&A, 423, 189
- Gratton R., 2000, in Pallavicini R., Micela G., Sciortino S., eds, ASP Conf. Ser. 198, Stellar Clusters and Associations: Convection, Rotation, and Dynamos, San Francisco, p. 225
- Grocholski A.J., Sarajedini A., 2003, MNRAS, 345, 1015
- Janes K.A., Tilley C., Lyngå G., 1988, AJ, 95, 771
- Jørgensen B.R., Lindegren L., 2005, A&A, 436, 127
- Karatas Y., Bilir S., Schuster W.J., 2005, MNRAS, 360, 1345
- Kharchenko N.V., Piskunov A.E., Röser S., Schilbach E., Scholz R.-D., 2005, A&A, 438, 1163
- Lata S., Pandey A.K., Sagar R., Mohan V., 2002, A&A, 388, 158
- Lebreton Y., 2000, ARA&A, 38, 35
- Loktin A.V., Gerasimenko T.P., Malisheva L.K., 2001, Astron. Astrophys. Trans., 20, 607
- Lyngå G., 1987, Catalogue of Open Cluster Data, 5th edition, CDS, Strasbourg
- Malysheva L.K., 1997, Astronomy Letters, 23, 585
- Mermilliod J.-C., 1981, A&A, 97, 235
- Paunzen E., Maitzen H.M., 2002, A&A, 385, 867
- Pena J.H., Peniche R., 1994, RMxAA, 28, 139
- Piatti A.E., Clariá J.J., Bica E., 1998, ApJS, 116, 263
- Pinsonneault M.H., Terndrup D.M., Hanson R.B., Stauffer J.R., 2004, ApJ, 600, 946
- Rees D.G., 1987, Foundations of Statistics, Chapman & Hall, London
- Schaerer D., Meynet G., Maeder A., Schaller G., 1993, A&AS, 98, 523
- Tadross A.L., 2001, New Astronomy, 6, 293
- Twarog B.A., Ashman K.M., Anthony-Twarog B.J., 1997, AJ, 114, 2556
- Winkler H., 1997, MNRAS, 287, 481